

## Choosing lime – what you need to know to make better decisions

Lisa Miller

*Southern Farming Systems (SFS)*

### Key messages

- Lime quality is affected mainly by its neutralising value and particle size and solubility plays a minor role affecting performance in the first 6 to 12 months
- A high quality lime has a NV above 90% and contains a high proportion of particle sizes less than 0.1 mm
- There are calculators available that can help growers compare the cost effectiveness of limes

### Background

The purchase of lime is a large upfront cost for growers and understandably they want to feel confident that their choice of lime is a good one and will ameliorate any acidity. Lime quality is a major factor that affects the extent of pH change and the time frame in which pH is changed.

There are three factors which control lime performance:

- Chemical composition determines the quantity of base applied to the soil and is accounted for in neutralising value (NV).
- Particle size determines the surface area of lime exposed to the soil volume and its distribution throughout the soil.
- Solubility determines the tendency of the material to dissolve and hence the rate of the reaction of a given particle size in the first 6 to 12 months.

### Lime Quality Indicators

NV is a measure of the lime's ability to neutralise acidity and is therefore the most important factor in selecting a lime. Pure calcium carbonate (or pure limestone) is taken as the standard with an NV of 100%. The higher the NV the more pure the product is. Lime products sold in Victoria commonly range in NV from 65% to 90%. Dolomite and burnt lime can have a NV greater than 100 percent due to their different chemical composition which is lighter in molecular weight compared with calcium carbonate but it won't necessarily neutralise more acid.

Particle size determines the amount of soil contact and coverage and how fast this reaction will occur. For example Cregan et al, 1989 mathematically calculated the amount of lime needed at different sizes to give complete coverage over one hectare. Particle diameters of less than 0.15 mm achieved complete coverage at rates close to 2.5 t/ha but 18 t/ha would be needed if the lime is greater than 1 mm (see table 1).

**Table 1.** Total amount of surface applied lime required to cover 1 ha at different particle diameters (mm).

Particle Diameter (mm)	Amount of lime needed to cover 1 ha (t/ha)
1	18.1
0.5	9.1
0.25	4.4
0.15	2.7

The effect of fineness on the speed of pH change is demonstrated in figure 1. This is based on an experiment at Wagga Wagga Agricultural Institute where a hard calcitic lime was incorporated to a 10 cm depth. This study is currently unpublished but was used by the NSW Department of Primary Industries 'Limes and Lemons' extension programs in the nineties.




Figure 1. The change in pH over time using different lime particle sizes.

The diagram shows that the finer the lime the greater the increase in pH and that coarse lime (2-5 mm) did not achieve the same pH change after 7 years. The bottom line is that coarse particles may never catch up. This is because the soil surrounding the lime particle reacts with the acidity and the pH becomes high and the lime ceases to dissolve, meanwhile the surrounding soil continues to acidify. Trials have also demonstrated that decreasing the particle size of lime will increase the rate of lime leaching which is important if trying to ameliorate subsurface acidity (Whitten, 2000). The implication is that the purchase of cheap coarse lime regardless of its NV may not be a good investment.

The emphasis on the importance of lime particle size has been lost over the years. Part of this reason is that the reporting of many quality factors including NV, particle fraction sizes and Effective Neutralising Value (ENV) was removed from mandatory reporting under the Victorian Fertiliser and Veterinary Regulations (Control of Use) in 2005. Now commonly only the proportions of particle sizes greater than 1 mm are reported. However, if you want a fast response to lime then a high percentage of the lime should be less than 0.1 mm. To make informed decisions on the purchase of lime, growers need to pay for the cost of a lime quality report by a NATA accredited testing laboratory which costs \$75 to \$100 per sample as there can be limited information on particle size provided to growers by lime producers.

Sometimes claims are made that particle size is not important in softer rock limes but this is not true. Particle size is important quality factor in all types of lime for soil coverage and for achieving pH change within a reasonable time frame. However solubility does vary amongst different lime types, although lime is generally regarded as being insoluble.

Solubility affects the lime's initial performance (first year) and for this reason is less important for consideration compared to NV and particle size which govern the limes overall performance. Solubility is linked to the type of liming material. NSW Agriculture field testing of 12 nationally available commercial limes found soft limes performed 20% more efficiently and dolomites 15% less efficiently than what their particle size model for calcitic hard limes predicted in the first 6 to 12 months after liming (Conyers et. al, 1995). By 12 months after application these differences disappeared.

Approximately 90% of liming products sold in Victoria are regarded as soft limes which are physically softer than harder limes. Examples of hard calcitic limes in Victoria are limes from Lilydale and Buchan and they are

generally finer than soft limes as they undergo more processing. Dolomite is a naturally occurring blend of magnesium carbonate and calcium carbonate.

### Tools to help growers compare limes

There are a number of tools that allow producers to make comparisons between different liming materials by accounting for both NV and particle size such as the ENV calculation commonly used in Victorian and South Australia shown below.

$$\text{ENV} = A + B + C$$

Where A = %  $\geq 0.85$  mm  $\times 0.1 \times \text{NV}\%$

Where B = %  $> 0.30$  mm but  $< 0.85$  mm  $\times 0.6 \times \text{NV}\%$

Where C = %  $< 0.30$  mm  $\times \text{NV}\%$

Generally the tools are based on an NV multiplied by the percentage of lime contained in different sieved fractions which are rated by their ability at changing pH within 12 months. All of the states agree that the finer the lime the higher its effectiveness at changing pH but at what particle size to set the 100% effectiveness rating varies across states. A comparison of effectiveness ratings is shown in table 2.

An ENV calculation first originated in US (Adams, 1971) where 0.3 mm particle size was set at 100% effective because a 0.3 mm sieve was the finest used. Lime comparisons using the ENV calculation were studied in pot trials by the Victorian Department of Agriculture in 1987 (Greenhill et al, 1987). They slightly changed the range of sieved particle fractions to fit their standard sieving practices. The calculators originating from WA use ratings modified from Cregan's DPI, NSW work in 1989 (Gazey, 2010). The NSW DPI calculator bases its effectiveness on field data from 1992 trials (Scott et al, 1992) and involved commercially available lime products. It best reflects that the finer the lime, the more effective it is. Its use is described in "Limes and Lemons".

Recent testing of limes by SFS showed a flaw in the particle size ratings used in the ENV calculation. It doesn't have the range in particle sizes to properly compare limes and should be revised. For example 4 out of 15 limes tested had greater than 10% of their lime exceeding 5 mm in size. Particles sizes of 5 mm shouldn't be rated the same reactivity as particle sizes of 1 mm. Particles of 5 mm could take 20 years to breakdown and result in poor pH change.

Table 2. Effectiveness ratings of different particle sizes which are used in different calculations and calculators for lime quality comparisons.

Vic. ENV calculation	NSW DPI Lime comparison calculator, 2003	WA DAFF calculator & the Soil Quality online Lime Comparison Calculator
< 0.3 mm = 100%	< 0.075 mm = 100%	<0.125 mm =100%
0.3 to 0.85 mm = 60%	0.075 to 0.15 mm = 58%	0.125 to 0.25 mm =100%
>0.85 mm = 10%	0.15 to 0.25 mm = 42%	0.25 to 0.5 mm = 100%
	0.5 to 1 mm = 34%	0.5 to 1 mm = 50%
	1 to 2 mm = 22%	> 1mm = 20%
	> 2 mm = 12%	

The cost effectiveness of different limes spread can be compared using LimeCheque a calculator developed by the South Australian government. This calculator is available from the SFS website under soil acidity.

### Acknowledgments

GRDC and CCMA through the support of DAFF. Brendan Scott, Research Scientist, Charles Sturt University for assistance in understanding this topic.

### **References**

Adams F, (1971) Soil, crop and economic benefits from lime use. In "Increasing lime use in the Tennessee Valley region," Bulletin Y-20 National Fertiliser Development Centre, Tennessee Valley Authority, Muscle Shoals, Alabama

Cregan PD, Hirth JR and Conyers MK (1989) Amelioration of soil acidity by liming and other amendments. In: Soil acidity and plant growth. Ed A D Robson. Academic Press, Australia.

Conyers MK, Scott B J, Fisher R and Lill W (1995) Predicting the field performance of twelve commercial liming materials from southern Australia. *Fertiliser Research* **44**, 151-161.

Gazey C (2010). Soil Acidity - A guide for WA farmers and consultants 2nd edition, Dept. of Agriculture and Food.

Greenhill N, Palmer B and Peverill, K, 1987. The reactivity of Victorian Liming materials in soils. State Chemistry Laboratory. Unpublished.

NSW Agriculture "Limes and Lemons" Unpublished.

Scott B J, Conyers M K, Fisher R and Lill W (1992) Particle size determines the efficiency of calcitic limestone in amending acidic soil. *Australian Journal of Agriculture Research* **43**, 1175-1185

Whitten M (2000) PhD thesis. Amelioration and treatment of agriculturally generated subsurface acidity in sandy soils in Western Australia.